

SPAY & NEUTER CENTER OF SOUTHERN NEVADA

EAST: 3475 E. Flamingo Rd. # 500 | WEST: 3210 S. Decatur Blvd. # 150 PHONE: 702-240-7729 (SPAY) · E-MAIL: Info@SpayNeuterLV.com www.SpayNeuterLV.com

Vaccination guidelines

We follow the guidelines set forth by AAHA (American Animal Hospital Association 2011) and The American Association of Feline Practitioners Feline Vaccine Advisory Panel

Canine

<u>DA2PPV</u> (5-in1) Distemper, Adenovirus I & II (hepatitis), Parainfluenza (part of the kennel cough complex) & Parvovirus. Vaccine is given every 3-4 weeks until 16 weeks of age. Booster in one year then once every 3 years. If over 16 weeks of age, give two doses 3-4 weeks apart. Booster in 1 year then once every 3 years. Vaccine protocols will vary depending on age and previous vaccine history. Speak with one of our staff members to determine current recommendations.

<u>Bordetella</u>- Kennel cough. Usually given intra-nasal but in some dogs given by injection. If less than 16 weeks of age, give 2 doses 3-4 weeks apart. Dogs over 16 weeks that have not received this vaccination will require 2 doses as well. Booster once a year. May give every 6 mos. depending on exposure (i.e. dogs that go to groomer, boarding, etc.).

<u>Rabies</u>- Can be given as early as 12 weeks, but usually given closer to 16 weeks. Then booster in one year. After that, it is given every 3 years (in NV). Same applies for an adult dog receiving rabies vaccination for the first time.

Feline:

FVRCP (3-in-1) Feline viral Rhinotracheitis, Calici-Virus & Panleukopenia (distemper). This is considered the distemper/upper respiratory vaccine. Vaccine is given every 3-4 weeks until 16 weeks of age, booster in one year then once every 3 years. If pet is over 16 weeks, give 2 vaccines 3-4 weeks apart, booster in one year then once every 3 years.

<u>FeLV</u>- Set of 2 vaccines 3-4 weeks apart in kittens or adults. Usually given at 12 week and 16 week visit. Then ONLY as exposure warrants (outdoor cat or if exposed to a lot of cats). It is recommended to test for FeLV/FIV before vaccination.

<u>Rabies</u>- Can be given as early as 12 weeks of age, but usually given closer to 16 weeks. Booster in one year then once every 3 years (in NV). Same applies for an adult cat receiving rabies vaccination for the first time.

These are basic guidelines. Certain pets may have different requirements based on exposure, health concerns and other factors. Please consult with our staff if you have specific questions or concerns.